

**Article Navigation**

FEBRUARY 14 2022

Autopsy Histopathologic Cardiac Findings in 2 Adolescents Following the Second COVID-19 Vaccine Dose James R. Gill, MD ; Randy Tashjian, MD; Emily Duncanson, MD*Arch Pathol Lab Med* (2022) 146 (8): 925–929.<https://doi.org/10.5858/arpa.2021-0435-SA> **Article history** **Context.—**

Myocarditis in adolescents has been diagnosed clinically following the administration of the second dose of an mRNA vaccine for coronavirus disease 2019 (COVID-19).

Objective.—

To examine the autopsy microscopic cardiac findings in adolescent deaths that occurred shortly following administration of the second Pfizer-BioNTech COVID-19 dose to determine if the myocarditis described in these instances has the typical histopathology of myocarditis.

Design.—

Clinical and autopsy investigation of 2 teenage boys who died shortly following administration of the second Pfizer-BioNTech COVID-19 dose.

Results.—

The microscopic examination revealed features resembling a catecholamine-induced injury, not typical myocarditis pathology.

Conclusions.—

The myocardial injury seen in these postvaccine hearts is different from typical myocarditis and has an appearance most closely resembling a catecholamine-mediated stress (toxic) cardiomyopathy. Understanding that these instances are different from typical myocarditis and that cytokine storm has a known feedback loop with catecholamines may help guide screening and therapy.

[Skip to Main Content](#)

myocarditis, although low, has been shown to increase after the receipt of the BNT162b2 vaccine, particularly after the second dose among young male recipients.¹¹ In addition, the first week after the second vaccine dose was found to be the main risk window.¹¹ The clinical presentation of myocarditis after vaccination was usually mild.¹¹

We report the autopsy results, including microscopic myocardial findings, of 2 teenage boys who died within the first week after receiving the second Pfizer-BioNTech COVID-19 dose. The microscopic findings are not the alterations seen with typical myocarditis. This suggest a role for cytokine storm, which may occur with an excessive inflammatory response, as there also is a feedback loop between catecholamines and cytokines.¹²

MATERIALS AND METHODS

The Connecticut Office of the Chief Medical Examiner and the Michigan Institute of Forensic Science and Medicine investigate all unexpected and unnatural deaths in their respective jurisdictions: Connecticut and the Michigan counties of Alcona, Gladwin, Huron, Lapeer, Ogemaw, and Saginaw.

Standard medicolegal autopsies were performed including gross, microscopic, and toxicologic testing. SARS-CoV-2 nasal swab testing was performed by reverse transcriptase–polymerase chain reaction assay. Tissues were sent to the National Center for Emerging and Zoonotic Infectious Diseases branch of the Centers for Disease Control and Prevention for molecular studies.

Cardiac molecular testing with sequence analysis and deletion/duplication testing of the 100 genes listed in Invitae's arrhythmia and cardiomyopathy comprehensive panel was performed.

RESULTS

The results of autopsies for 2 teenage boys who were found dead in bed 3 and 4 days after receiving the second dose of the Pfizer-BioNTech COVID-19 vaccine are presented ([Table](#)). Both boys were pronounced dead at home without attempted resuscitation.

Summary of Clinical and Autopsy Findings


Teenage boy B, BMI = 30 with obesity	520 g with biventricular dilatation and marked pulmonary edema (combined lung weight = 1481 g)	<p>histiocytic infiltrate</p> <p>PCR tissue testing performed by the CDC on heart and lung found no molecular evidence of SARS-CoV-2 infection</p> <p>Molecular testing on postmortem blood detected 2 variants of uncertain significance: <i>DOLK</i> (c.1257C.G [p.Ile419Met] heterozygous) and <i>MAP2K2</i> (c.581-3C>T [intronic] heterozygous)</p> <p>There was global myocardial injury similar to that seen above, but with more widespread transmural ischemic changes and more interstitial inflammation, again with a predominant neutrophil component with histiocytes and scant lymphocytes (Figures 5 through 7; Supplemental Figures 3 and 4). Several sections had transmural, confluent areas of hyper eosinophilic myocytes; confluent areas of contraction bands apart from any inflammation; and florid neutrophilic inflammation with some histiocytes. In this case, a subepicardial distribution of injury was not seen. There were no acute or organizing thrombi. PCR tissue testing performed by the CDC on heart and lung found no molecular evidence of SARS-CoV-2 infection</p>
--------------------------------------	--	---

Abbreviations: BMI, body mass index; CDC, Centers for Disease Control and Prevention; PCR, polymerase chain reaction.

View large

Boy A complained of a headache and gastric upset but felt better by postvaccine day 3. There was no history of prior medical problems (he took prescribed amphetamine/dextroamphetamine during the school year for attention deficit hyperactivity disorder but was not currently receiving it) or prior SARS-CoV-2 infection. Boy B had no complaints, prior health issues, or prior SARS-CoV-2 infection. Neither boy complained of fever, chest pain, palpitations, or dyspnea. The autopsies were unremarkable except for obesity in one boy and the cardiac findings (Figures 1 through 7; Supplemental Figures 1 through 4 [see supplemental digital content at <https://meridian.allenpress.com/aplm> in the August 2022 table of contents]). Unique cardiac findings in boy A included myocardial fibrosis and in boy B cardiac hypertrophy. There were no rashes or lymphadenopathy.

Figure 1



Skip to Main Content




View large

Download slide

Figure 3. Case A, heart: subepicardial fibrosis. This appears older than the timing of the first vaccine dose. This is a possible arrhythmogenic cardiomyopathy, but its appearance is more consistent with healed ischemia or inflammation (hematoxylin-eosin, original magnification $\times 40$).

Figure 4. Case A, heart: confluent areas of ischemia with contraction bands and coagulative myocytolysis (hematoxylin-eosin, original magnification $\times 200$).

Figure 5



Case B, heart: hypereosinophilic myocytes, contraction band necrosis, and coagulative myocytolysis. Inset: the infiltrate is predominantly neutrophilic (hematoxylin-eosin, original magnifications ×100 and ×400 [inset]).

Figure 6. Case B, heart: subepicardial coagulative myocytolysis/contraction band necrosis (hematoxylin-eosin, original magnification ×100).

Figure 7. Case B, heart: perivascular inflammation (hematoxylin-eosin, original magnification ×200).

Expanded forensic toxicologic testing was negative for medications and drugs of abuse. SARS-CoV-2 was not detected by postmortem swab (reverse transcriptase–polymerase chain reaction assay) in either boy. Cardiac sections were submitted from the right and left ventricles (12 sections in boy A and 29 sections in boy B). The cardiac conduction systems were not examined.

DISCUSSION

Myocarditis is an inflammatory disease of the myocardium, which may occur in isolation or as part of multiorgan/systemic immune-mediated disorders or reactions to exogenous/endogenous substances.¹³ The etiologies are varied and include infectious and noninfectious causes. Noninfectious causes include immune/autoimmune conditions (autoantigens, association with immune-mediated diseases, alloantigens, and allergens), drugs/toxic substances (eg, hypersensitivity or direct toxic effects), and other causes (eg, radiation, insect stings, snake bites).¹³ Lymphocytic myocarditis is the commonest histologic subtype, characterized by an inflammatory myocardial infiltrate typically comprising mononuclear cells. In the acute/active phases, it is usually accompanied by myocyte damage/necrosis.¹³ Although criteria are evolving, the Dallas criteria require “inflammatory infiltrates of the myocardium with necrosis and/or degeneration of *adjacent* myocytes, not typical of ischemic damage associated with coronary artery disease.”¹⁴⁻¹⁶

Toxic myocarditis is an etiologic classification involving direct myocardial injury by various drugs or substances.^{13,17,18} Although variable, the histologic features consist of 2 main patterns: an early stage with foci of solely necrotic/damaged myocytes and the later phase of “myocarditis.” Toxic myocarditis usually indicates inflammatory stages of catecholamine-induced myocardial injury. Catecholamine toxicity on the heart was first described in patients with pheochromocytoma.¹⁹⁻²¹ These lesions have been described in patients with subarachnoid hemorrhages and, more recently, in donor hearts rejected for transplantation in persons declared dead by neurologic criteria, secondary to catecholamine release during the “sympathetic storm” following brain death or administered as pharmacologic support (see [supplemental material](#)).^{22,23} The wide spectrum of these lesions has been studied in detail in routine pathology examination of donor hearts unsuitable for

transplantation.²²
[Skip to Main Content](#)

Both teenage boys had similar clinical presentations with no obvious cardiac symptoms. Their histopathology did not demonstrate a typical myocarditis. In those instances, one sees lymphocytic (or giant cell) infiltrates with

can develop in patients with extreme physical, chemical, or sometimes emotional stressors.²⁴⁻³¹

Stress cardiomyopathy is a catecholamine-mediated ischemic process seen in high catecholamine states in the absence of coronary artery disease or spasm.^{17,31} It has also been called “neurogenic myocardial injury” and “broken heart syndrome.”^{18,24-36} Surges in catecholamines may have several triggers (fight/flight response, adrenal pathology, etc). Proposed mechanisms for catecholamine-mediated stunning in stress cardiomyopathy include epicardial spasm, microvascular dysfunction, hyperdynamic contractility with midventricular or outflow tract obstruction, and direct effects of catecholamines on cardiomyocytes.³³

Catecholamine-mediated myocardial stunning may be due to direct myocyte injury, as elevated catecholamines decrease the viability of myocytes through cyclic adenosine monophosphate-mediated calcium overload. Catecholamines also are a potential source of oxygen-derived free radicals, which can interfere with sodium and calcium transporters, possibly resulting in myocyte dysfunction through increased transsarcolemmal calcium influx and cellular calcium overload.³⁷

Histologically, catecholamine effects have been associated with contraction band necrosis, characterized by hypercontracted sarcomeres, dense eosinophilic transverse bands, and an interstitial mononuclear inflammatory response that is distinct from the polymorphonuclear inflammation seen with infarction. In addition, the mononuclear cells are not causing the myocyte necrosis; there is a distinct, separate distribution.³⁷

We suspect that the acute cardiac changes seen in these 2 boys are the result of epinephrine-mediated effects on cardiomyocytes. These occurrences generally have a favorable prognosis; however, some patients may die from the underlying (noncardiac) cause of the myocardial findings (eg, as with subarachnoid hemorrhage). Histologically, diffuse hypereosinophilic myocytes, contraction bands, and coagulative myocytolysis are seen, with a patchy and random pattern and a neutrophilic/mononuclear cell infiltrate. With longer survival, global myocardial ischemia may develop.³⁷

This postvaccine reaction may represent an overly exuberant immune response, with the myocardial injury mediated by similar immune mechanisms to those described with SARS-CoV-2 and multisystem inflammatory syndrome cytokine storms.³⁸ Multisystem inflammatory syndrome is a rare systemic illness presenting with persistent fever and extreme inflammation following exposure to SARS-CoV-2. Affected children have persistent fever and may have acute abdominal pain with diarrhea or vomiting, muscle pain/malaise, and hypotension. Other reported symptoms include rashes, enlarged lymph nodes, and swelling.

A hypersensitivity reaction is in the differential diagnosis; however, infrequency or lack of eosinophils would be unusual. The common denominator of a hypersensitivity reaction is the eosinophilic infiltrate, which may be the major inflammatory component or part of a complex picture of mixed inflammation with lymphocytes, macrophages, plasma cells, poorly formed microgranulomas, and giant cells.³⁹ An autopsy study of 69 cases of

Despite a molecular investigation, the etiology of the fibrosis in case A is unclear. It is conceivable that this process first started with the first vaccination dose and the initial myocardial effects resolved and healed over time. The second dose may have restarted the process. One might expect some scarring/repair after a few weeks, although the scarring in case A appears more organized than the 3-week interval between the vaccine doses. Also, it is only in one of the cases. It remains possible that the fibrosis represents arrhythmogenic cardiomyopathy.

Unfortunately, cardiac molecular testing was equivocal.

Regardless of the etiology of the fibrosis, the extent of scarring by itself is potentially arrhythmogenic and may be a contributing factor with the acute postvaccine myocardial injury. Similarly, the cardiac hypertrophy in case B may have made the heart more susceptible to an arrhythmia. The key point is that since these boys died suddenly and unexpectedly in their sleep without resuscitation, if the arrhythmia had been due to the myocardial scar (boy A) or cardiomegaly (boy B), then the fulminant, global myocardial injury would not be an expected finding. These 2 clinical histories support the etiology of the acute myocardial injury as a primary factor, not a secondary agonal or postresuscitative artifact.

Two adults (ages 42 and 45 years) with myocarditis diagnosed histologically (one at autopsy and one by biopsy) following SARS-CoV-2 mRNA vaccinations were recently reported.⁴¹ One occurred 10 days after receiving the first Pfizer-BioNTech COVID-19 vaccine dose and the other occurred 14 days after receiving the second mRNA-1273 (Moderna) dose. Histologically, both were described as “fulminant” myocarditis with “multifocal cardiomyocyte damage associated with mixed inflammatory infiltration.” In addition to areas of myocyte necrosis associated with the inflammatory infiltrate, the photomicrographs demonstrate ischemic changes distinct from the inflammation, similar to our findings.

Cytokine storm has been described with an excessive and uncontrolled inflammatory response, and there is a feedback loop between catecholamines and cytokines.¹² Clinical complications may include cardiac compromise, respiratory distress, and hypercoagulation.⁴² The myocardial injury seen in these postvaccine hearts has a similar histologic appearance to catecholamine-mediated stress cardiomyopathy and severe SARS-CoV-2 infection, including myocarditis, which is associated with cytokine release syndrome.³⁸ Recognition that these instances are different from typical myocarditis and that cytokine storm has a known feedback loop with catecholamines may help guide screening, diagnosis, and therapy.

References

1. Wise J. Covid-19: should we be worried about reports of myocarditis and pericarditis after mRNA vaccines?

BMJ 2021; 373:n1635.

[Skip to Main Content](#)

[Google Scholar](#)

vaccine recipients: update from the Advisory Committee on Immunization Practices—United States, June 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70(27):977–982.

[Google Scholar](#)

4. Rosner CM, Genovese L, Tehrani BN, et al Myocarditis temporally associated with COVID-19 Vaccination. *Circulation.* 2021;144(6):502–505.

[Google Scholar](#)

5. Navar AM, McNally E, Yancy CW, O'Gara PT, Bonow RO. Temporal associations between immunization with the COVID-19 mRNA vaccines and myocarditis: the vaccine safety surveillance system is working. *JAMA Cardiol.* 2021;6(10):1117–1118.

[Google Scholar](#)

6. Muthukumar A, Narasimhan M, Li QZ, et al In-depth evaluation of a case of presumed myocarditis after the second dose of COVID-19 mRNA vaccine. *Circulation.* 2021;144(6):487–498.

[Google Scholar](#)

7. Montgomery J, Ryan M, Engler R, et al Myocarditis following immunization with mRNA COVID-19 vaccines in members of the US military. *JAMA Cardiol.* 2021;6(10):1202–1206.

[Google Scholar](#)

8. Marshall M, Ferguson ID, Lewis P, et al Symptomatic acute myocarditis in 7 adolescents after Pfizer-BioNTech COVID-19 vaccination. *Pediatrics.* 2021;148(3):e2021052478.

[Google Scholar](#)

9. Halushka MK, Vander Heide RS. Myocarditis is rare in COVID-19 autopsies: cardiovascular findings across 277 postmortem examinations. *Cardiovasc Pathol.* 2021;50: 107300.

[Google Scholar](#)

10. Fox SE, Falgout L, Vander Heide RS. COVID-19 myocarditis: quantitative analysis of the inflammatory infiltrate and a proposed mechanism. *Cardiovasc Pathol.* 2021;54: 107361.

[Google Scholar](#)

11. Mevorach D, Anis E, Cedar N, et al Myocarditis after BNT162b2 mRNA vaccine against Covid-19 in Israel. *N Engl J Med.* 2021;385(23):2140–2149.

[Google Scholar](#)

12. Staedtke V, Bai RY, Kim K, et al Disruption of a self-amplifying catecholamine loop reduces cytokine release syndrome. *Nature.* 2018;564(7735):273–277.

[Google Scholar](#)

13. Leone O, Pieroni M, Rapezzi C, Olivotto I. The spectrum of myocarditis: from pathology to the clinics. *Virchows Arch.* 2019;475(3):279–301.

[Google Scholar](#)

[Skip to Main Content](#)

14. Aretz HT. Myocarditis: the Dallas criteria. *Hum Pathol.* 1987;18(6):619–624.

[Google Scholar](#)

2013;18(6):673–681.

[Google Scholar](#)

17. Cebelin M, Hirsch C. Human stress cardiomyopathy: myocardial lesions in victims of homicidal assaults without internal injuries. *Hum Pathol.* 1980;11(2):123–132.

[Google Scholar](#)

18. Berman M, Ali A, Ashley E, et al Is stress cardiomyopathy the underlying cause of ventricular dysfunction associated with brain death? *J Heart Lung Transplant.* 2010;29(9):957–965.

[Google Scholar](#)

19. Wang AR, Dean SA, Grebe SK, Hood IC. Fatal catecholamine-induced cardiotoxicity associated with pheochromocytoma: report of a postpartum case and review of the literature. *Acad Forensic Pathol.* 2016;6(2):315–324.

[Google Scholar](#)

20. Frustaci A, Loperfido F, Gentiloni N, Caldarulo M, Morgante E, Russo MA. Catecholamine-induced cardiomyopathy in multiple endocrine neoplasia: a histologic, ultrastructural, and biochemical study. *Chest.* 1991;99(2):382–385.

[Google Scholar](#)

21. Van Vliet PD, Burchell HB, Titus JL. Focal myocarditis associated with pheochromocytoma. *N Engl J Med.* 1966;274(20):1102–1108.

[Google Scholar](#)

22. Leone O, Angelini A, Bruneval P, Potena L. *The Pathology of Cardiac Transplantation: A Clinical and Pathological Perspective.* Cham, Switzerland: Springer; 2016.

23. Pine DS, Tierney L, Jr. Clinical problem-solving: a stressful interaction. *N Engl J Med.* 1996;334(23):1530–1534.

[Google Scholar](#)

24. Lindsay J, Paixao A, Chao T, Pichard AD. Pathogenesis of the Takotsubo syndrome: a unifying hypothesis. *Am J Cardiol.* 2010;106(9):1360–1363.

[Google Scholar](#)

25. Indorato F, Bartoloni G. Post-mortem Takotsubo cardiomyopathy diagnosis: the challenge is open! *Forensic Sci Med Pathol.* 2016;12(2):227–228.

[Google Scholar](#)

26. Angelini P. Do pathologists agree on how to diagnose Takotsubo cardiomyopathy? *Forensic Sci Med Pathol.* 2016;12(2):226.

[Google Scholar](#)

27. Indorato F, Akashi YJ, Rossitto C, Raffino C, Bartoloni G. Takotsubo cardiomyopathy associated with rupture of the left ventricular apex: assessment of histopathological features of a fatal case and literature

[Google Scholar](#)

29. Savall F, Dedouit F, Telmon N, Rouge D. Takotsubo cardiomyopathy in a jeweler following a hold-up: forensic considerations. *J Forensic Sci.* 2013;58(6):1648–1650.

[Google Scholar](#)

30. Toni C, Iannaccone F, Chella P, Basolo F, Pucci A. Sudden death in a case of recurrent Takotsubo syndrome. *Forensic Sci Med Pathol.* 2019;15(4):595–597.

[Google Scholar](#)

31. Akashi YJ, Goldstein DS, Barbaro G, Ueyama T. Takotsubo cardiomyopathy: a new form of acute, reversible heart failure. *Circulation.* 2008;118(25):2754–2762.

[Google Scholar](#)

32. Assuncao LA, Grams AC, Ribeiro CS, Magalhaes T. “Broken heart” syndrome in an elder abuse case: forensic considerations from the case. *Forensic Sci Med Pathol.* 2011;7(2):217–221.

[Google Scholar](#)

33. Lyon AR, Rees PS, Prasad S, Poole-Wilson PA, Harding SE. Stress (Takotsubo) cardiomyopathy—a novel pathophysiological hypothesis to explain catecholamine-induced acute myocardial stunning. *Nat Clin Pract Cardiovasc Med.* 2008;5(1):22–29.

[Google Scholar](#)

34. Boland TA, Lee VH, Bleck TP. Stress-induced cardiomyopathy. *Crit Care Med.* 2015;43(3):686–693.

[Google Scholar](#)

35. Akashi YJ, Nef HM, Mollmann H, Ueyama T. Stress cardiomyopathy. *Annu Rev Med.* 2010;61: 271–286.

[Google Scholar](#)

36. Sacha J, Maselko J, Wester A, Szudrowicz Z, Pluta W. Left ventricular apical rupture caused by Takotsubo cardiomyopathy—comprehensive pathological heart investigation. *Circ J.* 2007;71: 982–985.

[Google Scholar](#)

37. Wittstein IS, Thiemann DR, Lima JA, et al Neurohumoral features of myocardial stunning due to sudden emotional stress. *N Engl J Med.* 2005;352(6):539–548.

[Google Scholar](#)

38. Gubbi S, Nazari MA, Taieb D, Klubo-Gwiezdzinska J, Pacak K. Catecholamine physiology and its implications in patients with COVID-19. *Lancet Diabetes Endocrinol.* 2020;8(12):978–986.

[Google Scholar](#)

39. Al Ali AM, Straatman LP, Allard MF, Ignaszewski AP. Eosinophilic myocarditis: case series and review of literature. *Can J Cardiol.* 2006;22(14):1233–1237.

[Google Scholar](#)[Skip to Main Content](#)

40. Burke AP, Saenger J, Mullick F, Virmani R. Hypersensitivity myocarditis. *Arch Pathol Lab Med.* 1991;115(8):764–769.

[Google Scholar](#)

2-related multisystem inflammatory syndrome in children: a systematic review. *J Pediatric Infect Dis Soc.* 2020;9(6):746–751.

[Google Scholar](#)

Author notes

Supplemental digital content is available for this article at <https://meridian.allenpress.com/aplm> in the August 2022 table of contents.

The authors have no relevant financial interest in the products or companies described in this article.

Supplementary data

[arpa-146-06-04_s01](#)- pdf file



ORCHARD
Software

Call us today to learn more about Orchard's cutting edge laboratory information solutions.

LEARN MORE

(800) 856-1948
www.orchardsoft.com

Access to industry specific jobs! The career center is a cohesive destination for CAP members and nonmembers to connect with prospective employers and employees in the laboratory industry.

Head and Neck Pathologist
Houston, TX
**University of Texas MD
Anderson Cancer Center**

**Pathologist - Stevens Point,
Wisconsin**
Stevens Point, WI
Aspirus Health

Academic Hematopathologist
Phoenix, AZ
Mayo Clinic

**Talent Pool - Forensic
Pathologist (Fee for Service)**
Toronto, ON
**Ministry of the Solicitor
General**

VIEW JOBS

Sponsored Video

How to Perform a Run and...



View Metrics

CITING ARTICLES VIA

Web Of Science (18)

Google Scholar

Clinicopathologic and Molecular Characterization of High-Risk Human Papillomavirus–Positive Carcinomas of the Urinary Tract

Neslihan Kayraklioglu, MD, PhD, Bradley A. Stohr, MD, PhD, Emily Chan, MD, PhD

Perivascular Epithelioid Cell–Family Tumors in Children, Adolescents and Young

Adults: Clinicopathologic Features in 70 Cases :

Phoebe M. Hammer, MD, Angus Toland, MD, Muhammad Shaheen, MD, Archana Shenoy, MD, Ashwini ESNakula, MD, MS, M. John Hicks, MD, PhD, DDS, Mikako Warran, MD, Alyaa Al-Ibraheemi, MD, Jessica L. Davis, MD, Serena Y. Tan, MD

Digital and Computational Pathology Are Pathologists' Physician Extenders

Casey P. Schukow, DO, Timothy Craig Allen, MD, JD

Continued Positive Job Search Experience for New Pathologists Seeking First Employment During the COVID-19 Pandemic (2020–2022)

Julie Katz Karp, MD, Amanda Lofgreen, MS, Kristen Johnson, PhD, Melissa R. George, DO, Stephanie Barak, MD, Bronwyn H. Bryant, MD, Mary P. Berg, MD, John M. Childs, MD, Barbara E. C. Knollman-Ritschel, MD, Victor G. Prieto, MD, PhD, Kristie L. White, MD, Cindy B. McCloskey, MD

Contributions From the University of Michigan 2022 New Frontiers in Pathology Conference

Maria Westerhoff, MD, Lauren B. Smith, MD, Liron Pantanowitz, MD, PhD, MHA

Reply to “Fellowship Board Pass Rates Rising: Analysis of Pathology Subspecialty Examination Performance”

Gary W. Procop, MD, Ritu Nayar, MD

Nonspecificity of Immunohistochemistry for Mycobacteria Species Using a Rabbit Polyclonal Antibody

Justin T. Kelley, MD, MPH, Lauren Kroll-Wheeler, MD, Steven Hrycaj, PhD, May P. Chan, MD, Jerome Cheng, MD, Laura W. Lamps, MD

Steroidogenic Acute Regulatory Protein Is a Useful Marker for Sex-Cord-Stroma Tumors and Normal and Neoplastic Adrenocortical Tissue

Maximilian Lennartz, MD, Daniela Amezada, Doris Höflmayer, MD, Sebastian Dwertmann Rico, MD, Clara von Barga, MD, Simon Kind, MD, Viktor Reiswich, MD, Florian Viehweger, MD, Florian Lutz, MD, Veit Bertram, MD, Christoph Fraune, MD, Natalia Gorbokon, MD, Sören

MUSCLE MESCHERHYMATOCYTIC TUMORS: A REVIEW OF THE

Current Knowledge

Pavel Dundr, PhD, Radoslav Matěj, PhD, Jan Hojný, PhD, Nikola Hájková, PhD, Kristýna Němejcová, PhD, Michaela Kendall Bártů, PhD

Uncommon Tumors of the Lung: Recently Described and Rediscovered Tumors

Cesar A. Moran, MD

Get Email Alerts

Article Activity Alert

Publish Ahead of Print Alert

Latest Issue Alert